

# Confluence Detection

Wieger Wesselink

September 2019

## 1 Confluence

Consider the following untimed linear process specification  $P$ , with initial state  $d_0$ .

$$P(d) = \sum_{i \in I} \sum_{e_i} c_i(d, e_i) \rightarrow a_i(f_i(d, e_i)) \cdot P(g_i(d, e_i))$$

We distinguish different kinds of confluence. Let summand  $j \in I$  be the index of a  $\tau$ -summand, and  $i \in I$  be the index of an arbitrary summand. In the sequel we abbreviate  $c_i(d, e_i)$ ,  $f_i(d, e_i)$  and  $g_i(d, e_i)$  with  $c_i$ ,  $f_i$  and  $g_i$ . Note that for the  $\tau$ -summand with index  $j$  we have  $a_j = \tau$  and  $f_j = []$ .

### 1.1 Trivial confluence

Trivial confluence is defined as

$$C_{trivial}(i, j) = \forall d, e_i, e_j : (c_i \wedge c_j) \Rightarrow (a_i = \tau) \wedge (g_i = g_j)$$

Note that trivial confluence only applies to  $\tau$ -summands. In isolation it is not a very useful property to check.

### 1.2 Triangular confluence

Triangular confluence is defined as

$$C_{triangular}(i, j) = \forall d, e_i, e_j : (c_i \wedge c_j) \Rightarrow (c_i[d := g_j] \wedge (f_i = f_i[d := g_j]) \wedge (g_i[d := g_j] = g_i))$$

### 1.3 Commutative confluence

Commutative confluence is defined as

$$C_{commutative}(i, j) = C_{trivial}(i, j) \vee \forall d, e_i, e_j : (c_i \wedge c_j) \Rightarrow \exists e'_i, e'_j : \left( \begin{array}{l} c_i[d := g_j, e_i := e'_i] \\ \wedge c_j[d := g_i, e_j := e'_j] \\ \wedge (f_i = f_i[d := g_j, e_i := e'_i]) \\ \wedge (g_i[d := g_j, e_i := e'_i] = g_j[d := g_i, e_j := e'_j]) \end{array} \right) \quad (1)$$

The reason for adding the term  $C_{trivial}(i, j)$  is probably that otherwise a simple  $\tau$ -summand like

$$(n = 0) \rightarrow \tau \cdot P(n = 1)$$

is not even confluent with itself.

## 1.4 Square confluence

Square confluence is defined as

$$C_{square}(i, j) = C_{trivial}(i, j) \vee \forall d, e_i, e_j : (c_i \wedge c_j) \Rightarrow c_i[d := g_j] \wedge c_j[d := g_i] \wedge (f_i = f_i[d := g_j]) \wedge (g_i[d := g_j] = g_j[d := g_i])$$

It is obtained from  $C_{commutative}(i, j)$  by taking  $e'_i = e_i$  and  $e'_j = e_j$ .




Figure 1: square commutative confluence, with  $d_1 = g_j[d := g_i]$




Figure 2: triangular confluence




Figure 3: trivial confluence, with  $a_j = \tau$